arXiv:astro-ph/9710302v1 27 Oct 1997

Minkowski Functionals in Cosmology: An Overview

Jens Schmalzing
Ludwig–Maximilians–Universität, Theresienstraße 37, 80333 München, Germany
Max–Planck–Institut für Astrophysik, Karl–Schwarzschild–Straße 1,
85740 Garching, Germany

Minkowski functionals have recently been introduced into cosmology as novel tools for studying the large–scale distribution of matter in the Universe. We present a brief overview of the method, including its mathematical foundations as well as some completed and upcoming applications.

1 Mathematical background

Useful insights into the large–scale structure of the Universe are possible by investigating the morphological properties of density fields and point sets. Integral geometry[1,2] provides the mathematical framework for such considerations. Hadwiger's Theorem[3] states that in order to measure the morphology of a d–dimensional pattern, the $d+1$ Minkowski functionals are sufficient.

2 Application to point processes

Originally[4], applications of Minkowski functionals in cosmology focussed on the Boolean grain model of a point set given, for example, by the coordinates of galaxies in redshift space. The Boolean model constructs the union set of balls centered around the members of the point set. Their common radius can be employed as a diagnostic parameter, as the Minkowski functionals of the union set change in a characteristic fashion with increasing radius.

The comparison of a subsample of Abell and ACO clusters and several varieties of the Cold Dark Matter scenario[5] demonstrates the discriminative power of Minkowski functionals. Figure 1 clearly shows the failure of the Standard Cold Dark Matter model. Other Cold Dark Matter scenarios, such as Broken Scale Invariance or an additional cosmological constant, lead to much better agreement.

Significant fluctuations on large scales are found in an analysis of the IRAS 1.2Jy galaxy catalogue[6]. Since the results shown in Figure 2 for samples of $100h^{-1}$Mpc depth remain robust on even larger scales and under various tests, they are probably a feature of the large–scale matter distribution in the Universe, rather than an effect in the catalogue. An explanation, for example in terms of cosmic variance, is still lacking, and a topic of ongoing study.



Figure 1: This plot shows the Minkowski functionals for the Abell/ACO cluster catalogue (solid line) compared to several realizations of SCDM mock catalogues (shaded area).



Figure 2: Here we display the Minkowski functionals of volume–limited subsamples of $100h^{-1}$Mpc depth from the northern and southern hemispheres of the IRAS 1.2Jy galaxy catalogue.

3 Outlook on further applications

One of the Minkowski functionals is the Euler characteristic or genus, which has long been used in cosmology[7]. As an extension, all Minkowski functionals of a smoothed random field may be considered[8]. Furthermore, applications to cosmic microwave background anisotropy maps[9] are possible.

References

1. R. Schneider. *Convex bodies: the Brunn–Minkowski theory.* Cambridge 1993.
2. W. Weil. In *Convexity and its applications*, pp. 360–412. Basel 1983.
3. H. Hadwiger. *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie.* Berlin 1957.
4. K. R. Mecke, T. Buchert & H. Wagner. *A&A* **288** (1994) 697–704.
5. M. Kerscher et al. *MNRAS* **284** (1997) 73–84.
6. M. Kerscher, J. Schmalzing, T. Buchert & H. Wagner. *A&A* submitted, astro-ph/9704028, 1997.
7. J. R. Gott III, A. L. Melott & M. Dickinson. *ApJ* **306** (1986) 341–357.
8. J. Schmalzing & T. Buchert. *ApJ* **482** (1997) L1–L4.
9. J. Schmalzing & K. M. Górski. *MNRAS* submitted, astro-ph/9710185, 1997.